POST-SURGERY INSTRUCTIONS: LIMITED INCISION BROWLIFT

Patient Name	
--------------	--

Date

Surgery Date

Once your surgery is completed, you must follow all the instructions given to you in order to heal properly and have a good outcome.

The following instructions are your obligation. Use this as a checklist of progress as you heal. Included are normal post-surgical experiences and key health considerations that may be a cause of concern.

TYPICAL POST-OPERATIVE SYMPTOMS

Typical symptoms of brow lift surgery, and signs to watch for following brow lift surgery include the following:

Tightness in the scalp, forehead, and brow: Bruising and swelling in the eyelid region: These are normal experiences as the skin, tissues and sensory nerves heal. Pain medication will help you cope with any discomfort. **Consistent sharp pain should be reported to our office immediately.**

Asymmetry: the brows look different, or heal differently: The brows may look or feel quite different from one another in the days following surgery. This is normal; no face in nature or following surgery is perfectly symmetrical.

CALL THE OFFICE IMMEDIATELY IF YOU EXPERIENCE ANY OF THE FOLLOWING:

- A high fever (over 101°), severe nausea and vomiting, continued dizziness or incoherent behavior, such as hallucinations.
- Any pain that cannot be controlled by your pain medication.
- Bright red skin that is hot to the touch.
- Excessive bleeding or fluid seeping through the incisions.
- A severely misshapen eyelid or excessive bruising or fluid retention that is localized to one region.

To alleviate any discomfort, and to reduce swelling, you may apply cool (not cold) compresses to your scalp, forehead, and brows. Do not apply ice or anything frozen directly on the skin. Do not apply anything cool on your cheeks or neck. Soak soft plain white washcloths or gauze squares in ice water and wring out well. Apply directly to the forehead, brow and eyes, but do not apply any pressure. Apply cool compresses for no longer than 20-minute intervals.

COMPRESSION

You may be given a compression garment or wrap immediately following surgery. Wear this exactly as directed. Remove it only as directed for cleansing incisions or showering.

DAY OF SURGERY INSTRUCTIONS

You will only be released to the care of a responsible adult. All of these instructions must be clear to the adult who will monitor your health and support you, around the clock in the first 24 hours following surgery.

	impor	but not bed rest: While r ant is that you are ambula 10 minutes every 2 hours	tory, meaning that	at you are walking un	der your own strength.		
	Recline, do not lie down: This will be more comfortable for you, and can reduce swelling. Always keep your head elevated. Do not bend forward or over.						
	caffeir drinks	nutrition: Fluids are critic ne-free and green tea-free . You must consume at le bus food for the first 24 hou	beverages incluc ast 8 ounces of fl	ling fruit juices and w	ater, milk and yogurt		
		all medication, exactly as ations you must take inclu		al pain medication, ar	ntibiotics and other		
Antibiotic:				mg		x per day	
Pain medicat	ion:			mg		x per day	
Ointment							
Eye drops							
Other			<u> </u>				
Supplements	:						
	surger remov Do no follow	your incisions clean. You ry. A cotton swab soaked re any crusting near your s of smoke. Smoking can gro ing surgery. You must not . Do not engage in any str	in warm water is titches. Do not re eatly impair your smoke.	appropriate for clean emove any staples, s safety prior to surger	sing incisions. Do not utures or steri-strips. y and you ability to heal		

TWO TO SEVEN DAYS FOLLOWING SURGERY

During this time you will progress with each day that passes. Ease into your daily activities. You will receive clearance to begin driving or return to work at your post-operative visit, or within:

days

Your post-operative visit is scheduled for:

• Continue to cleanse wounds as directed; you may shower and shampoo your hair. Take a warm, not hot shower. Do not rub your scalp or your incisions. Use a mild shampoo and no styling products. If you must dry your hair, do so only with a hand-held hair dryer on the coolest setting and do not allow the compressed air to blow directly onto your incisions or your face.

- **Apply ointment and skincare as directed.** Do not use any glycolic, retinoid or other potentially irritating skincare products on your face.
- **Take antibiotic medications and supplements as directed.** Take pain medication only as needed. You may wish to switch from prescription pain medication to acetaminophen or ibuprofen.
- Continue to wear dark lens, large framed sunglasses and a wide-brimmed hat whenever you are outdoors. You may begin wearing reading glasses as soon as it is comfortable for you.
- Continue to keep you head elevated, including when sleeping.
- **Do not resume any exercise other than regular walking.** Walking is essential every day to prevent the formation of blood clots.
- Maintain a healthy diet. Do not smoke. Do not consume alcohol.

Post Operative treatment: Make certain to schedule all of the post-operative treatments as prescribed

ONE to FOUR WEEKS FOLLOWING SURGERY

As you resume your normal daily activities, you must continue proper care and healing.

- Continue your wound care as directed. Do not use any styling products or tools until all stitches are removed AND until your incisions no longer have any crusting or scabbing.
- Refrain from direct sun exposure. Continue to wear your sunglasses and a wide-brimmed hat. If you are
 outdoors, apply at least an SPF 30 at least 30 minutes prior to sun exposure. Your face is highly susceptible
 to sunburn or the formation of irregular, darkened pigmentation.
- **Do not smoke.** While incisions may have sealed, smoking deprives your body of necessary oxygen that can result in poorly healed, wide, raised scars.
- Refrain from any strenuous exercise and from bending or lifting.
- You may begin sleeping in a modified reclining position. However do not sleep lying flat or on your stomach. If you are a side sleeper, two pillows under your head, and a soft pillow under your mid-back and shoulders may offer more comfort.

Follow-up as directed. Your second post-operative visit is scheduled for:

SIX WEEKS FOLLOWING SURGERY

Healing will progress; swelling and bruising continue to diminish.

- You may ease into your regular fitness routine. However, protective eyewear and a hat are necessary when outdoors.
- Discomfort or tightness and tingling in your forehead, brow and scalp will resolve.
- No need to resume smoking. You have now gone 10 weeks (4 weeks prior to surgery and 6 weeks following) without a cigarette. For your long-term health, there is no need to resume smoking.

• You may notice some hair loss at the incision sites. This is normal. If it becomes excessive, or the hair is shedding in clumps, please notify our office.

YOUR FIRST YEAR

- Continue good skincare and sun protection, healthy nutrition and fitness.
- Schedule any complementary procedures, as recommended. Botulinum injections or other treatments may be recommended to enhance your results, and to help your results to be long-lasting.
- Your scars will continue to refine. If they become raised, red or thickened, or appear to widen, contact our office. Early intervention is important to achieving well-healed scars. Scars are generally refined to fine incision lines one year after surgery.
- A one-year post surgery follow-up is recommended. However you may call our office at any time with your concerns or for needed follow-up.

Your appearance will change with age. Your facial appearance may change too. You may wish to undergo revisional surgery at a later date to help maintain your appearance throughout life. Contact our office with any of your questions or concerns, at any time.

I have read and understand all of the above instructions. I understand that following these instructions is solely my responsibility. I understand that it is also my responsibility to ask my doctor and his or her staff any questions I have related to these instructions or about my procedure, health and healing.

Patient Signature

Date

Printed Name of Patient

Signature of Practice Representative and Witness