POST-SURGERY INSTRUCTIONS: NECKLIFT

Patient Name	Date	
	0 0.	
	Surgery Date	

Once your surgery is completed, you must follow all the instructions given to you in order to heal properly and have a good outcome.

The following instructions are your obligation. Use this as a checklist of progress as you heal. Included are normal post-surgical experiences and key health considerations that may be a cause of concern.

TYPICAL POST-OPERATIVE SYMPTOMS

Typical symptoms of neck lift surgery; and signs to watch for following a neck lift include:

Tightness or numbness of chin, jaw and neck. Stiffness of the neck: Bruising and swelling of the neck or a pale, swollen complexion. These are normal experiences as the skin, tissues and sensory nerves heal. Pain medication and muscle relaxants will help you cope with any discomfort. Consistent sharp pain should be reported to our office immediately.

If you have a drain placed in any incision, you may also experience localized discomfort at the drain site.

CALL THE OFFICE IMMEDIATELY IF YOU EXPERIENCE ANY OF THE FOLLOWING:

- A high fever, (over 101°) severe nausea and vomiting, continued dizziness or incoherent behavior, such as hallucinations.
- Any pain that cannot be controlled by your pain medication.
- Bright red skin that is hot to the touch.
- Excessive bleeding or fluid seeping through the incisions.
- A severely misshapen appearance, excessive bruising or fluid retention that is localized to one region.

To alleviate any discomfort, and to reduce swelling, you may apply cool (not cold) compresses to your jaw region, not to your neck. Do not apply ice or anything frozen directly on the skin. Soak soft plain white washcloths or gauze squares in ice water and wring out well. Apply directly to the jaw, but do not apply any pressure. Apply cool compresses for no longer than 20-minute intervals. Do not apply any heat.

COMPRESSION

You may be placed in a compression garment or wrap immediately following surgery. Wear this exactly as directed. Remove it only as directed for cleansing incisions or showering.

DAY OF SURGERY INSTRUCTIONS

You will only be released to the care of a responsible adult. All of these instructions must be clear to the adult who will monitor your health and support you around the clock in the first 24 hours following surgery.

	importa	ant is that you are	While rest is imperambulatory: meany 2 hours engaged	ning that you a	re walking under	your own strength	J .
	Recline, do not lie down: This will be more comfortable for you, and can reduce swelling. Always keep your head elevated. Do not bend forward or over.						
	caffein drinks.	e-free and green	tea-free beverage ime at least 8 ound	s including fruit	juices and wate		,
		II medication, extitions you must to	xactly as prescritake include:	oed. Oral pain r	medication, antib	iotics and other	
Antibiotic:	_				mg		x per day
Pain medica	tion:				mg		x per day
Ointment: Other:	_						
Supplements	s: _						
	surgery remove you ha	 A cotton swab any crusting nea ve any drains pla 	soaked in warm war your stitches. Doced, follow the ins	vater is appropr o not remove a tructions for cle	iate for cleansing any staples, sutu eansing and carir	d for a short time a g incisions. Do no res or steri-strips. ng for drains. nd your ability to h	t If
	Relax.	Do not engage	in any stressful ac	tivities. Take c	are of no one. Le	et others tend to yo	ou.
TWO TO SE	VEN DA	YS FOLLOWING	SURGERY				
			each day that pas our post-operative		your daily activi days	ties. You will rece	ive clearance
Your	post-op	erative visit is sch	neduled for:				

- Continue to cleanse wounds as directed; you may shower and shampoo your hair. Take a warm, not hot shower. Do not rub your scalp or your incisions. Use a mild shampoo and no styling products if you have incisions within your scalp. If you must dry your hair, do so only with a hand-held hair dryer on the coolest setting and do not allow the compressed air to blow directly onto your incisions or your neck.
- Apply ointment and skincare as directed. Do not use any glycolic, retinoid or other potentially irritating skincare products on your face until you receive clearance to do so.

- Take antibiotic medications and supplements as directed. Take pain medication only as needed. You
 may wish to switch from prescription pain medication to acetaminophen or ibuprofen.
- Avoid sun exposure of any kind to your face and neck. If you must be outdoors for any reason, apply
 at least an SPF 30 sunscreen on your face and neck, and wear a wide brimmed hat or cover your neck
 with a scarf or other wrap.
- Continue to keep your head elevated, including when sleeping.
- Do not resume any exercise other than regular walking. Walking is essential every day to prevent the formation of blood clots.
- Maintain a healthy diet. Do not smoke. Do not consume alcohol.

Post-operative treatment: prescribed:	Make certain to schedule all of the post-operative treatments as
	□

ONE TO FOUR WEEKS FOLLOWING SURGERY

As you resume your normal daily activities, you must continue proper care and healing.

- Continue your wound care as directed. If your incisions are within the scalp: Do not use any styling products or tools until all stitches are removed AND until your incisions no longer have any crusting or scabbing. If your incisions are on the skin: Do not use any make-up until the stitches are removed AND until your incisions no longer have any crusting or scabbing.
- Refrain from direct sun exposure. Continue to wear your sunglasses and a wide-brimmed hat. If you are
 outdoors, apply at least an SPF 30 at least 30 minutes prior to sun exposure. Your face is highly susceptible
 to sunburn or the formation of irregular, darkened pigmentation.
- **Do not smoke.** While incisions may have healed, smoking deprives your body of necessary oxygen that can result in poorly healed, wide, raised scars.
- Refrain from any strenuous exercise and from bending or lifting.
- You may begin sleeping in a modified reclining position. If you are a side sleeper, two pillows under your head and a soft pillow under your mid-back and shoulders may offer more comfort.

SIX WEEKS FOLLOWING SURGERY

Healing will progress; swelling and bruising continue to diminish.

- You may ease into your regular fitness routine.
- Discomfort or tightness and tingling in your face will resolve.
- **No need to resume smoking.** You have now gone 10 weeks (4 weeks prior to surgery and 6 weeks following) without a cigarette. For your long-term health, there is no need to resume smoking.
- You may notice some hair loss at incision sites within the scalp. This is normal. If it becomes excessive, or the hair is shedding in clumps, please notify our office.

YOUR FIRST YEAR

- Continue good skincare and sun protection, healthy nutrition and fitness.
- Schedule any complementary procedures, as recommended. Botulinum injections or specific skincare treatments may be recommended to enhance your results, and to help your results to be long-lasting.
- Your scars will continue to refine. If they become raised, red or thickened, or appear to widen, contact our
 office. Early intervention is important to achieving well-healed scars. Scars are generally refined to fine
 incision lines one year after surgery.
- A one-year post surgery follow-up is recommended. However you may call our office at any time with your concerns or for needed follow-up.

Your appearance will change with age. Your facial appearance may change too. You may wish to undergo revision surgery at a later date to help maintain your appearance throughout life. Contact our office with any of your questions or concerns, at any time.

I have read and understand all of the above instructions. I understand that following these instructions is solely my responsibility. I understand that it is also my responsibility to ask my doctor and his or her staff any questions I have related to these instructions or about my procedure, health and healing.				
Patient Signature	Date			
Printed Name of Patient				
	Circulation of Departing Department of the condition and Milling			
	Signature of Practice Representative and Witness			