### POST-SURGERY INSTRUCTIONS: MINI ABDOMINOPLASTY

Patient Name	Date	
	Surgery Date	

Once your surgery is completed, you must follow all the instructions given to you in order to heal properly and have a good outcome.

The following instructions are your obligation. Use this as a checklist of progress as you heal. Included are normal post-surgical experiences and key health considerations that may be a cause of concern.

#### TYPICAL POST-OPERATIVE SYMPTOMS

Typical symptoms of mini abdominoplasty and signs to watch for following limited tummy tuck surgery include the following:

**Tightness and stiffness in abdomen: Bruising, swelling and redness. Tingling, burning or intermittent shooting pain:** These are normal experiences as the skin, tissues and sensory nerves heal. Pain medication and muscle relaxants will help you cope with any discomfort. If you have drains, you may experience additional localized discomfort. **Consistent sharp pain should be reported to our office immediately.** 

Skin firmness, hypersensitivity or lack of sensitivity: This is normal and will gradually resolve over time.

**Shiny skin or any itchy feeling:** Swelling can cause the skin in treated areas to appear shiny. As the healing process advances, you may also find a mild to severe itchy feeling. An antihistamine like Benadryl can help to alleviate severe, constant itchiness. **If the skin becomes red and hot to the touch, contact our office immediately.** 

### CALL THE OFFICE IMMEDIATELY IF YOU EXPERIENCE ANY OF THE FOLLOWING:

- A high fever, (over 101°) severe nausea and vomiting, continued dizziness or incoherent behavior, such as hallucinations.
- Any pain that cannot be controlled by your pain medication.
- Bright red skin that is hot to the touch.
- Excessive bleeding or fluid seeping through the incisions.
- A severely misshapen region anywhere on your abdomen or bruising that is localized to one specific point of the abdomen.

To alleviate any discomfort, and to reduce swelling, you may apply cool, not cold compresses to the treated region. Crushed ice or ice packs must be wrapped in a towel before being applied to the skin. Do not apply ice or anything frozen directly to the skin. Apply cool compresses, for no longer than 20-minute intervals.

# **DAY OF SURGERY INSTRUCTIONS**

Whether you are released after surgery or after an overnight stay in a recovery center or hospital, you will only be released to the care of a responsible adult. All of these instructions must be clear to the adult who will monitor your health and support you, around the clock in the first 24 hours following surgery.

	importa Spend	ant is that you a 10 minutes eve	re ambulatory, meanir ry 2 hours engaged in	ant in the early stages of hing that you are walking und light walking indoors as you comfortable for you, and	der your own strength. ou recover.
	caffein drinks.	e-free and greei	n tea-free beverages i sume at least 8 ounces	surgery. Stick to non-carbo ncluding fruit juices and was s of fluid every 2 hours. Sti	ater, milk and yogurt
		III medication, of ations you must		d. Oral pain medication, ar	ntibiotics and other
Antibiotic:	_			mg	x per day
Pain medica	ition:			mg	x per day
Ointment: Eye drops: Other:	-				x per day
Supplement	s: _				
	time af for clea ointme compr	ter surgery. Keansing incisions on the sterions wraps the instructions of the sterions of the instructions of the sterions of	ep dressings clean an  Do not remove any sestrips, then apply 4x4  over the gauze. If yo	sions will seep fluid and so d dry. A cotton swab with steri-strips over your stitch gauze pads over your inc the have a drain placed in your ord drained fluid on the <b>Dr</b> a	peroxide is appropriate es. Apply anti-bacterial isions. <b>Replace any</b> our incisions, carefully
				the clock. Follow the insteanse your incision or to e	
			ng can greatly impair must not smoke.	your safety prior to surgery	y and your ability to heal
		Do not engagene, and let othe		ities. Do not lift , push or p	pull anything. Take care

## TWO TO SEVEN DAYS FOLLOWING SURGERY

During this time you will progress as each day passe	s. Ease into your daily activities.	You will receive clearance to
begin driving or return to work at your post-operative	visit, or within	days.
	-	
Your post-operative visit is scheduled for:		

- Continue to clean your wounds as directed; you may shower. Take a warm, not hot shower. Do not take
  a bath. Limit your shower to 10 minutes. Avoid getting your incisions wet. Do not remove any steri-strips. Do
  not rub your incisions. Apply a fragrance free moisturizer to the surrounding skin, however not on your
  incisions.
- Take antibiotic medications and supplements as directed. Take pain medication and muscle relaxants only as needed. You may wish to switch from prescriptive pain medication to acetaminophen or ibuprofen.
- Continue to wear your compression garment around the clock.
- Begin scheduled lymphatic drainage and massage therapy. This can help soften any firmness or contour irregularities.
- Do not resume any exercise other than regular walking. Walking is essential every day to prevent the formation of blood clots.
- No sun exposure. If you plan to go outdoors for any reason, wear protective clothing. Avoid any direct sun exposure.
- Maintain a healthy diet. Do not smoke. Do not consume alcohol.

# ONE to FOUR WEEKS FOLLOWING SURGERY

As you resume your normal daily activities, you must continue proper care and healing.

- Continue your wound care as directed.
- Refrain from weight-bearing or abdominal crunching exercises. Continue walking. A daily, brisk 20-minute walk is recommended.
- **Do not smoke.** While incisions may have sealed, smoking deprives your body of necessary oxygen that can result in poorly healed, wide, raised scars.
- Continue to wear your compression garment as directed.
- Continue to attend massage therapy as scheduled.
- **Practice good sun protection.** Do not expose your abdomen to direct sunlight. If you are outdoors, apply at least an SPF 30 at least 30 minutes prior to sun exposure and wear protective clothing. The skin of your abdomen is highly susceptible to sunburn or the formation of irregular, darkened pigmentation.

Follow-up as directed.	Your second post-operative visit is scheduled for:	
		<u> </u>

## SIX WEEKS FOLLOWING SURGERY

Healing will progress and your lower body settles into a more final shape and position.

- You may ease into your regular fitness routine. However realize that your body may require some time to return to previous strength.
- Discomfort or tightness and tingling of the skin will resolve.
- No need to resume smoking. You have now gone 10 weeks (4 weeks prior to surgery and 6 weeks following) without a cigarette. For your long-term health, there is no need to resume smoking.

#### YOUR FIRST YEAR

- Continue healthy nutrition, fitness and sun protection.
- Your scars will continue to refine. If they become raised, red or thickened, or appear to widen, contact our
  office. Early intervention is important to achieving well-healed scars. Scars are generally refined to fine
  incision lines one year after surgery.
- A one-year post surgery follow-up is recommended. However you may call our office at any time with your concerns or for needed follow-up.

Your body will change with age. The appearance of your abdomen will change too. Although the outcomes of a limited tummy tuck are generally permanent, any significant weight gain or loss, pregnancy as well as the normal influences of aging can cause changes to your appearance. You may wish to undergo revision surgery at a later date to help maintain your appearance throughout life. Contact our office with any of your questions or concerns, at any time.

I have read and understand all of the above instructions. I understand that following these instructions is solely my responsibility. I understand that it is also my responsibility to ask my doctor and his or her staff any

questions I have related to these instructions or about	my procedure, health and healing.
Patient Signature	Date
Printed Name of Patient	-

Signature of Practice Representative and Witness