POST-SURGERY INSTRUCTIONS: SCAR REVISION SURGICAL

Patient Name		
Patient Name		

Date _____

Surgery Date

Once your scar revision is completed, you must follow all the instructions given to you in order to heal properly and have a good outcome.

The following instructions are your obligation. Use this as a checklist of progress as you heal. Included are normal post-surgical experiences and key health considerations that may be a cause of concern.

TYPICAL POST-OPERATIVE SYMPTOMS

Typical symptoms and signs to watch for following surgical scar revision include the following:

Tingling, burning, redness, tightness at the surgical site. These are normal experiences as the skin, tissues and sensory nerves heal. Pain medication will help you cope with any discomfort. **Consistent sharp pain should be reported to our office immediately.** You may also feel minor discomfort at the graft donor site, if you had your own skin grafted to revise the scar.

Shiny skin or any itchy feeling: Swelling can cause the skin to appear shiny. As the healing process advances, you may also find a mild to severe itchy feeling at the scar revision or any graft donor site. An antihistamine like Benadryl can help to alleviate severe, constant itchiness. If the skin becomes red and hot to the touch, contact our office immediately.

CALL THE OFFICE IMMEDIATELY IF YOU EXPERIENCE ANY OF THE FOLLOWING:

- A high fever, (over 101°) severe nausea and vomiting, continued dizziness or incoherent behavior, such as hallucinations.
- Any pain that cannot be controlled by your pain medication.
- Bright red skin that is hot to the touch.
- Excessive bleeding or fluid seeping through the incisions.
- A severely misshapen or swollen appearance at the wound site.

To alleviate any discomfort, and to reduce swelling, you may apply cool, not cold compresses to the wound site. Crushed ice or ice packs must be wrapped in a towel before being applied to the skin. Do not apply ice or anything frozen directly to the skin. Apply cool compresses for no longer than 20-minute intervals.

DAY OF SURGERY INSTRUCTIONS

If you have had sedation of any kind, you will only be released to the care of a responsible adult. All of these instructions must be clear to the adult who will monitor your health and support you, around the clock in the first 24 hours following surgery.

	import	but not bed rest. While rest is ant is that you are ambulatory, I I 10 minutes every 2 hours enga	meaning that y	ou are walking under	your own strength.					
	Keep the wound site elevated. If the scar revision is of the face or upper body, a reclined position is best. If your scar revision is of the lower body, keep your legs elevated.									
	caffeir drinks	nutrition. Fluids are critical fol ne-free, and green tea-free beve . You must consume at least 8 vith soft, bland, nutritious food fo	erages includin ounces of fluid	g fruit juices and wate every 2 hours. If you	r, milk, and yogurt					
		all medication, exactly as pres ations you must take include:	scribed : Oral բ	pain medication, antib	iotics and other					
Antibiotic	-			mg	x per day					
Pain medication			mg	x per day						
	-			mg	x per day					
Other	-									
Supplements										
	seep f	ge your incision dressings. Y luid and some blood for a short swab with peroxide is appropria over your stitches. Apply anti-b pad.	time after surg ate for cleansir	ery. Keep dressings ng incisions. Do not re	clean and dry. A emove any steri-					
	wear,	compression as directed. If y you must wear this around the c car revision can result in the for	clock, as direct	ed. If you do not follow						
		t smoke. Smoking can greatly ng surgery. You must not smol		fety prior to surgery a	nd your ability to heal					
		. Do not engage in any stressfu in in the region of your scar revi		o not do anything that	can stress or stretch					

TWO TO SEVEN DAYS FOLLOWING SURGERY

During this time you will progress with each day that passes. Ease into your daily activities. You will receive clearance to begin driving or return to work at your post-operative visit, or within:

days

Your post-operative visit is scheduled for:

- Continue to cleanse wounds as directed; you may shower. Take a warm, not hot shower. Do not take a bath. Limit your shower to 10 minutes. Do not remove any steri-strips. Do not rub your incisions. Apply ointment and any compression as directed.
- Take medications and supplements as directed. Pain medication such as acetaminophen or ibuprofen should ease your discomfort.
- **Proper sun protection is vital.** You must not allow direct or indirect sun exposure to your wounds. Wear protective clothing, a wide-brimmed hat and at least an SPF 30 if you must be outdoors.
- **Do not resume any exercise other than regular walking.** Walking is essential every day to prevent the formation of blood clots.
- Maintain a healthy diet. Do not smoke. Do not consume alcohol.

ONE TO FOUR WEEKS FOLLOWING SURGERY

As you resume your normal daily activities, you must continue proper care and healing.

- Discomfort or tightness and tingling will resolve.
- **Continue wound care as directed.** This includes any ointment, compression tape or other instructions you have been given.
- Ease into your fitness routine. Avoid aerobic exercise that may stress or stretch the skin in the area of your scar revision.
- **Do not smoke.** While incisions may have sealed, smoking deprives your body of necessary oxygen that can result in poorly healed, wide, raised scars. For your long-term health, there is no need to resume smoking.
- **Practice good sun protection.** Do not expose your skin to direct sunlight. If you are outdoors, apply at least an SPF 30 at least 30 minutes prior to sun exposure and wear protective clothing. Your skin is highly susceptible to sunburn or the formation or irregular, darkened pigmentation.

Follow-up as directed. Your second post-operative visit is scheduled for:

YOUR FIRST YEAR

- Continue healthy nutrition, fitness, and sun protection.
- Your scars will continue to refine. If they become raised, red or thickened, or appear to widen, contact our office. Early intervention is important to achieving well-healed scars. Scars are generally refined to fine incision lines one year after surgery.
- A one-year post surgery follow-up is recommended. However, you may call our office at any time with your concerns or for needed follow-up.

Your skin and scars may change with age. Contact our office with any of your questions or concerns, at any time.

I have read and understand all of the above instructions. I understand that following these instructions is solely my responsibility. I understand that it is also my responsibility to ask my doctor and his or her staff any questions I have related to these instructions or about my procedure, health and healing.

Patient Signature

Date

Printed Name of Patient